



Test yourself

district levels and strategic deployment of human resources (trichiasis surgeons), equipment, and consumables.

The prevalence of active disease has decreased to levels below the threshold recommendation for district-level mass drug administration (MDA), and so the programme has been able to stop this activity in 84 % of the districts where trachoma is present (Figures 1 and 2). This has been due to the high annual rates of coverage with Zithromax® and tetracycline during MDA, strong data collection efforts, and conducting surveys to assess impact.

To address the F and E components, the PNLC conducted several different activities at the same time. These were:

- training for a variety of community groups and leaders (local women's groups, religious/village leaders, and community volunteers) in trachoma prevention
- broadcasting of health messages on community radio stations
- development of a trachoma school health curriculum that is being taught in primary schools
- household latrine construction and community-led total sanitation. Since 2009, PNLC support has assisted in the construction of 53,090 latrines.

Future plans

The PNLC and partners will continue to build upon the gains made over the past 5 years and support the planning and implementation of SAFE strategy activities. The national programme is refining its surgical planning in order to reach the remaining 27,000 people estimated to need trichiasis surgery, thereby achieving the 'elimination goal' of less than one case of trichiasis per 1,000 persons.

Simultaneously, MDA to reduce transmission of trachoma will continue in communities where the prevalence remains high. Surveillance will also continue in areas where MDA has stopped. Social mobilisation and community sensitisation through radios, community volunteers, and women's groups will play a vital role in supporting attitudes and behaviours that help prevent the transmission of disease, strengthen disease knowledge, and decrease the number of people who refuse treatment or surgery. Ongoing latrine construction will continue to provide household access to safe disposal of faeces.

With thanks to Sanoussi Bamani, Seydou Goita, Yaya Kamissoko, Sadi Moussa, Sidi Coulibaly, Aryc W. Mosher, and Emily Toubali for their contributions to this article.

Test your understanding of the concepts covered in this issue and discuss any points of interest with your manager or a colleague. *Produced in collaboration with the International Council of Ophthalmology (ICO).*

1. Think about 'balancing the books' and sustainability		True	False
a	External donor funds are best used for training and capacity building	<input type="checkbox"/>	<input type="checkbox"/>
b	Governments are not responsible for paying for eye care	<input type="checkbox"/>	<input type="checkbox"/>
c	If you charge for services, everyone should pay the same	<input type="checkbox"/>	<input type="checkbox"/>
d	Pharmacy and spectacle sales are two areas where income can be generated	<input type="checkbox"/>	<input type="checkbox"/>
2. Think about patient flow, accounting and procurement		True	False
a	Buying smaller quantities of consumables, more frequently, saves money	<input type="checkbox"/>	<input type="checkbox"/>
b	Intraocular lenses (IOLs) should be on the procurement list if a hospital offers cataract surgery	<input type="checkbox"/>	<input type="checkbox"/>
c	You don't need to have a computer to set up an accounting system	<input type="checkbox"/>	<input type="checkbox"/>
d	Reducing the number of times a patient must visit the hospital, e.g. for cataract surgery, saves costs	<input type="checkbox"/>	<input type="checkbox"/>

ANSWERS

1a. True, b. False, c. True, d. True. 2a. False, b. True, c. True, d. True. You can start on paper. 2a. False, b. True, c. True, d. True. Some patients will pay more for value-added services. 2b. True, c. True, d. True. Bulk buy medicines and consumables yearly or quarterly. 2c. False, d. True. Governments must cover some of the costs or organise health insurance, as in Ghana.


PICTURE QUIZ

Diagnose this

A ground-glass appearance of the cornea is noted immediately after cataract surgery (figure) and there is a +3 anterior chamber reaction. What condition do you suspect?

What is the most likely diagnosis?

- Endophthalmitis
- Mechanical trauma to the cornea
- Intraoperative introduction of a toxic substance into the eye
- Fuchs' corneal dystrophy



© American Academy of Ophthalmology, used by permission

ANSWER

Bacterial contamination of the cleaning bath detergent for surgical instruments may also cause acute corneal oedema following cataract surgery. Oedema in these cases can be abrupt, resulting in immediate corneal swelling. Toxic anterior segment syndrome (TASS) is an acute, sterile anterior segment inflammation following generally uneventful cataract and anterior segment surgery. Rapid onset of corneal oedema and absence of a hypopyon are the distinguishing factors in infectious endophthalmitis. Most patients with TASS will develop symptoms and signs within 12 to 24 hours of the operation.

Answer: Introduction of a toxic substance into the eye. Mechanical trauma to the endothelium during surgery is considered to be the most significant factor influencing postoperative corneal oedema; however, toxic substances may inadvertently be introduced into the eye, causing immediate corneal oedema. This may come from intraocular irrigation solutions or topical and intracameral anaesthetics. Intraocular medications that have resulted in corneal toxicity include epinephrine (now available preservative free), various preparations of lidocaine, benzalkonium chloride-preserved viscoelastic, vancomycin at doses greater than 1 mg/mL, and inadvertent exposure of the endothelium to 5% povidone-iodine.